

Erratum

As a result of a production error, several articles appearing in the Central Surgical Association (*Surgery*, 2007; Vol. 142, No. 4:433-644) and American Association of Endocrine Surgeons (*Surgery*, 2007; Vol. 142, No. 6:785-1030) special focus issues were published without their respective discussions. The articles affected have now been updated online to include the missing discussion material. *Surgery* apologizes to the authors for this significant oversight.

The following articles were affected:

Laparoscopic liver surgery for everyone: the hybrid method

Alan J. Koffron, Robert D. Kung, Gregory B. Auffenberg, and Michael M. Abecassis, *Chicago, Ill* (*Surgery*, 2007; Vol. 142, No. 4:463-8.)

Cellular edema regulates tissue capillary perfusion after hemorrhage resuscitation

El Rasheid Zakaria, MD, PhD, Na Li, MD, PhD, Paul J. Matheson, PhD, and Richard N. Garrison, MD, *Louisville, Ky* (*Surgery*, 2007; Vol. 142, No. 4:487-96.)

Experience with deceased donor kidney transplantation in 114 patients over age 60

Phillip S. Moore, MD, Alan C. Farney, MD, PhD, Erica L. Hartmann, MD, Jeffrey Rogers, MD, William Doares, PharmD, Michael D. Gautreaux, PhD, Samy S. Iskandar, MBBCh, PhD, Gloria Hairston, BS, Patricia L. Adams, MD, and Robert J. Stratta, MD, *Winston-Salem, NC* (*Surgery*, 2007; Vol. 142, No. 4:514-23.)

A contemporary analysis of outcomes for operative repair of type A aortic dissection in the United States

Brian S. Knipp, MD, G. Michael Deeb, MD, Richard L. Prager, MD, Candace Y. Williams, MD, Gilbert R. Upchurch, Jr., MD, and Himanshu J. Patel, MD, *Ann Arbor, Mich* (*Surgery*, 2007; Vol. 142, No. 4:524-8.)

Laparoscopy or conventional open surgery for patients with ileocolonic Crohn's disease? A prospective study

Alessandro Fichera, MD, Stephanie L. Peng, BA, Nicholas M. Elisseou, BS, Michele A. Rubin, RN, CSN, APN, and Roger D. Hurst, MD, *Chicago, Ill* (*Surgery*, 2007; Vol. 142, No. 4:566-71.)

Duodenum-preserving head resection for chronic pancreatitis: an institutional experience and national survey of usage

Thomas K. Varghese, MD, and Richard H. Bell, Jr., MD, *Chicago, Ill* (*Surgery*, 2007; Vol. 142, No. 4:588-93.)

Laparoscopic colorectal surgery is safe in the high-risk patient: a NSQIP risk-adjusted analysis

Carol A. McCloskey, MD, Mark A. Wilson, MD, PhD, FACS, Steven J. Hughes, MD, FACS, and George M. Eid, MD, FACS, *Pittsburgh, Pa* (*Surgery*, 2007; Vol. 142, No. 4:594-7.)

Eliminating respiratory intensive care unit stay after gastric bypass surgery

Peter T. Hallowell, MD, Thomas A. Stellato, MD, Marianna C. Petrozzi, MD, Margaret Schuster, RN, BSN, Kristen Graf, RN, BSN, Ann Robinson, BA, and John J. Jasper, MD, *Cleveland, Ohio* (*Surgery*, 2007; Vol. 142, No. 4:608-12.)

Fistulojejunostomy for the management of refractory pancreatic fistula

Rajalakshmi R. Nair, MD, Andrew M. Lowy, MD, Benjamin McIntyre, MD, Jeffrey J. Sussman, MD, Jeffrey B. Matthews, MD, and Syed A. Ahmad, MD, *Cincinnati, Ohio, and Chicago, Ill* (*Surgery*, 2007; Vol. 142, No. 4:636-42.)

Consequences of parathyroidectomy in patients with "mild" sporadic primary hyperparathyroidism

Denise M. Carneiro-Pla, MD, George L. Irvin, III, MD, FACS, and Herbert Chen, MD, *Miami, Fla, and Madison, Wis* (*Surgery*, 2007; Vol. 142, No. 4:795-9.)

Pheochromocytoma penetrance varies by RET mutation in MEN 2A

Frank J. Quayle, MD, Elizabeth A. Fialkowski, MD, Ronald Benveniste, MD, PhD, and Jeffrey F. Moley, MD, *St. Louis, Mo, and Houston, Tex* (*Surgery*, 2007; Vol. 142, No. 4:800-5.)

0039-6060/\$ - see front matter

© 2008 Mosby, Inc. All rights reserved.

doi:10.1016/j.surg.2007.12.001

Increased prevalence of breast cancer among patients with thyroid and parathyroid disease

Carolyn N. Garner, MD, Rebecca Ganetzky, BS, Jennifer Brainard, MD, Jeffrey P. Hammel, MS, Eren Berber, MD, Allan E. Siperstein, MD, and Mira Milas, MD, *Cleveland, Ohio*
(*Surgery*, 2007; Vol. 142, No. 4:806-13.)

Clinical, genetic and radiographic analysis of 108 patients with von Hippel-Lindau disease (VHL) manifested by pancreatic neuroendocrine neoplasms (PNENs)

Joseph A. Blansfield, MD, Lynda Choyke, DPM, Shane Y. Morita, MD, Peter L. Choyke, MD, James F. Pingpank, MD, H. Richard Alexander, MD, Geoffrey Seidel, RN, Yvonne Shutack, RN, Nargiza Yuldashaeva, RN, Michelle Eugeni, RN, David L. Bartlett, MD, Gladys M. Glenn, MD, PhD, Lindsay Middleton, RN, W. Marston Linehan, MD, and Steven K. Libutti, MD, *Bethesda, Md*
(*Surgery*, 2007; Vol. 142, No. 4:814-8.)

Long-term follow-up after parathyroidectomy for radiation-induced hyperparathyroidism

Giuseppe Ippolito, MD, F. Fausto Palazzo, MS, FRCS, Frederic Sebag, MD, and Jean François Henry, MD, *Marseille, France, and London, United Kingdom*
(*Surgery*, 2007; Vol. 142, No. 4:819-22.)

Trends in thyroid cancer demographics and surgical therapy in the United States

Ian Mitchell, MD, Edward H. Livingston, MD, Alice Y. Chang, MD, Shelby Holt, MD, William H. Snyder, III, MD, Ildiko Lingvay, MD, MPH, and Fiemu E. Nwariaku, MD, *Dallas, Tex*
(*Surgery*, 2007; Vol. 142, No. 4:823-8.)

Pancreatoduodenal surgery in patients with multiple endocrine neoplasia type 1: operative outcomes, long-term function, and quality of life

Y. Nancy You, MD, MHS, Geoffrey B. Thompson, MD, William F. Young, Jr, MD, Dirk Larson, MA, David R. Farley, MD, Melanie Richards, MD, and Clive S. Grant, MD, *Rochester, Minn*
(*Surgery*, 2007; Vol. 142, No. 4:829-36.)

The incidence of cancer and rate of false-negative cytology in thyroid nodules greater than or equal to 4 cm in size

Kelly L. McCoy, MD, Noel Jabbour, Jennifer B. Ogilvie, MD, N. Paul Ohori, MD, Sally E. Carty, MD, and John H. Yim, MD, *Pittsburgh, Pa*
(*Surgery*, 2007; Vol. 142, No. 4:837-44.)

Interpretation of intra-operative PTH changes in patients with multi-glandular primary hyperparathyroidism (pHPT)

Giorgos Constantine Karakousis, MD, Dale Han, MD, Rachel Rapaport Kelz, MD, Deepika Nemani, BS, Jagajan Karamacharya, MD, Robert Roses, MD, Phyllis A. Gimotty, PhD, and Douglas L. Fraker, MD, *Philadelphia, Pa*
(*Surgery*, 2007; Vol. 142, No. 4:845-50.)

Does the presence of additional thyroid nodules on ultrasound alter the risk of malignancy in patients with a follicular neoplasm of the thyroid?

Rebecca S. Sippel, MD, Dina M. Elaraj, MD, Elham Khanafshar, MD, Electron Kebebew, MD, Quan-Yang Duh, MD, and Orlo H. Clark, MD, *San Francisco, Calif*
(*Surgery*, 2007; Vol. 142, No. 4:815-7.)

Laryngoscopy in thyroid surgery—essential standard or unnecessary routine?

Katja Schlosser, MD, Mike Zeuner, MD, Maria Wagner, MD, Emily P. Slater, PhD, Emilio Domínguez Fernández, MD, Matthias Rothmund, MD, and Katja Maschuw, MD, *Marburg and Frankfurt/Main, Germany*
(*Surgery*, 2007; Vol. 142, No. 4:858-64.)

“Subclinical Cushing’s syndrome” is not subclinical: improvement after adrenalectomy in 9 patients

Ian C. Mitchell, MD, Richard J. Auchus, MD, PhD, Kavita Juneja, MD, Alice Y. Chang, MD, Shelby A. Holt, MD, William H. Snyder, III, MD, and Fiemu E. Nwariaku, MD, *Dallas, Tex*
(*Surgery*, 2007; Vol. 142, No. 4:900-5.)

Utilization of total thyroidectomy for papillary thyroid cancer in the United States

Karl Y. Bilimoria, MD, David J. Bentrem, MD, John G. Linn, MD, Andrew Freel, MD, Jen Jen Yeh, MD, Andrew K. Stewart, MA, David P. Winchester, MD, Clifford Y. Ko, MD, MS, MSHS, Mark S. Talamonti, MD, and Cord Sturgeon, MD, MS, *Chicago, Ill, Los Angeles, Calif, Chapel Hill, NC, and Evanston, Ill*
(*Surgery*, 2007; Vol. 142, No. 4:906-13.)

Functional parathyroid carcinoma: long-term treatment outcome and risk factor analysis

Masatoshi Iihara, MD, PhD, Takahiro Okamoto, MD, MSc, PhD, Rumi Suzuki, MD, Akiko Kawamata, MD, Toshio Nishikawa, MD, PhD, Makio Kobayashi, MD, PhD, and Takao Obara, MD, PhD, *Tokyo, Japan*
(*Surgery*, 2007; Vol. 142, No. 4:936-43.)

The preoperative exclusion of thyroid carcinoma in multinodular goiter: dynamic contrast-enhanced magnetic resonance imaging versus ultrasonography-guided fineneedle aspiration biopsy

Fatih Tunca, MD, Y. Giles, MD, A. Salmaslioglu, MD, A. Poyanli, MD, D. Yilmazbayhan, MD, T. Terzioglu, MD, and S. Tezelman, MD, *Istanbul, Turkey*
(*Surgery*, 2007; Vol. 142, No. 4:992-1002.)

Medullary thyroid microcarcinoma recommendations for treatment—A single-center experience

Christian Scheuba, MD, Klaus Kaserer, MD, Christian Bieglmayer, PhD, Reza Asari, MD, Phillip Riss, MD, Ralph Drosten, MD, and Bruno Niederle, MD, *Vienna, Austria*
(*Surgery*, 2007; Vol. 142, No. 4:1003-10.)